

+ BIOBRANE[◇]

Temporary Biosynthetic Wound Dressing

Biosynthetic dressing for use on extensive
body surface area wounds¹⁻³



Smith+Nephew

BIOBRANE[◇]
Temporary Biosynthetic
Skin Dressing

www.smith-nephew.com

BIOBRANE[®] Temporary Biosynthetic Wound Dressing has been successfully used on partial thickness burns of a patient with 85% Total Body Surface Area (TBSA)¹

Features and benefits



Conforms to surface irregularities^{1,4-6}

A flexible and conformable composite dressing that is easy to apply, and has adherent properties^{1,4-8}



Accelerates healing

BIOBRANE Dressing increased the speed of healing and epithelisation^{*9-12}



Transparent

BIOBRANE Dressing is transparent, allowing direct wound inspection^{4,6,13}



Paediatric use

BIOBRANE Dressing can be used in paediatric populations^{3,5,6,14,15}



Reduced healing time[†]

BIOBRANE Dressing reduced healing time,[†] which may lead to a quicker discharge^{9-11,16}

Indications

BIOBRANE Temporary Wound Dressing is indicated for:

- Covering clean partial thickness burn wounds
- Split thickness donor sites

BIOBRANE Glove is indicated for:

- Covering clean partial thickness burn wounds of the hand



*Compared to Silver Sulfadiazine (SSD) (BIOBRANE 9.5–13.7 days vs SSD 11.2-21.3 days)

†Compared with Silver Sulfadiazine (SSD) (BIOBRANE 9.5-13.7 days vs SSD 11.2-21.3 days)

BIOBRANE Temporary Wound Dressing

S+N Code	Size	Carton
66800046	13cm x 13cm	1
66800047	13cm x 38cm	1
66800048	25cm x 38cm	1
66800174	13cm x 13cm	5
66800175	13cm x 38cm	5
66800176	25cm x 28cm	5
66800177	38cm x 50cm	1

BIOBRANE Temporary Wound Glove

S+N Code	Size	Carton
66800049	Glove - Paediatric	1
66800051	Glove - Medium	1
66800052	Glove - Large	1
66800181	Glove - Paediatric	2
66800182	Glove - Small	2
66800183	Glove - Medium	2
66800184	Glove - Large	2

Advanced Wound Management
Smith & Nephew Medical Ltd
Croxley Park Building 5
Lakeside, Hatters Lane, Watford
Hertfordshire WD18 8YE - UK

www.smith-nephew.com

T +44 (0) 1482 225181
F +44 (0) 1482 328326

©Trademark of Smith+Nephew
All Trademarks acknowledged
©May 2022 Smith+Nephew
AWM-AWD-32780. GMC1481

For detailed product information, including indications for use, contraindications, precautions and warnings, please consult the product's applicable Instructions for Use (IFU) prior to use.

References

1. Erdman D, Hussman J, Kucan JO. Treatment of a severe alkali burn. *Burns*. 1996;22(2):141–146.
2. Lo CH, Akbarzadeh S, McLean C, et al. Wound healing after cultured epithelial autografting in patients with massive burn injury: a cohort study. *J Plast Reconstr Aesthet Surg*. 2019;72(3):427–437.
3. Muganza A, Cullingworth L. A Prospective Single-Centre Randomised Controlled Study to Compare the Time to Healing of Partial Thickness Burn Wounds Treated with Versajet, Biobrane and Acticoat Therapy. *Glob J Med Res*. 2014;14(6):5–12.
4. Farroha A, Frew Q, El-Muttardi N, Philp B, Dziewulski P. The Use of BIOBRANE[®] to Dress Split-thickness Skin Graft in Paediatric Burns. *Ann Burns Fire Disasters*. 2013;XXVI(2):94–97.
5. Fan C, Pek CH, Por YC, Lim GJS. Biobrane dressing for paediatric burns in Singapore: A retrospective review. *Singapore Med J*. 2018;59(7):360–365.
6. Leshner AP, Curry RH, Evans J, et al. Effectiveness of Biobrane for treatment of partial thickness burns in children. *J Pediatr Surg*. 2011;46(9):1759–1763.
7. Austin RE, Merchant N, Shahrokhi S, Jeschke MG. A comparison of Biobrane and cadaveric allograft for temporizing the acute burn wound: Cost and procedural time. *Burns*. 2015;41(4):749–753.
8. Yang J-Y, Tsai Y-C, Noordhoff MS. Clinical comparison of commercially available Biobrane preparations. *Burns*. 1989;15(3):197–203.
9. Barret JP, Dziewulski P, Ramzy PI, Wolf SE, Desai MH, Herndon DN. Biobrane versus 1% silver sulfadiazine in second-degree pediatric burns. *Plastic and Reconstructive Surgery*. 2000;105(1):62–65.
10. Gerding RL, Imbembo AL, Fratiante RB. Biosynthetic skin substitute vs. 1% silver sulfadiazine for treatment of inpatient partial-thickness thermal burns. *J Trauma*. 1988;28(8):1265–1269.
11. Wasiak J, Cleland H, Campbell F. Dressings for superficial and partial thickness burns (Review). *Cochrane Database Syst Rev*. 2008(4).
12. Kumar RJ, Kimble RM, Boots R, Pegg SP. Treatment of partial-thickness burns: a prospective, randomized trial using Transcyte. *ANZ J Surg*. 2004;74(8):622–626.
13. Lang EM, Eiberg CA, Brandis M, Stark GB. Biobrane in the treatment of burn and scald injuries in children. *Ann Plast Surg*. 2005;55(5):485–489.
14. Selvarajah D, Bolu BK, Harvey J, et al. Acticoat versus biobrane: a retrospective review on the treatment of paediatric mid-dermal torso burns. *Int J Burn Trauma*. 2019;9(4):82–87.
15. Cassidy C, St Peter SD, Lacey S, et al. Biobrane versus duoderm for the treatment of intermediate thickness burns in children: a prospective, randomized trial. *Burns*. 2005;31(7):890–893.
16. Lal S, Barrow RE, Wolf SE, et al. Biobrane improves wound healing in burned children without increased risk of infection. *Shock (Augusta, Ga)*. 2000;14(3):314–318; 318–319.