

# T.I.M.E. clinical decision support tool

## Assess patient, wellbeing and wound

Establish diagnosis and baseline characteristics for appropriate support and comorbidities that may impact healing. Record wound type, location, size, wound bed condition, signs of infection / inflammation, pain location and intensity, comorbidities, adherence / concordance to treatment


## Bring in multi-disciplinary team and informal carers to promote holistic patient care

Record referral to others such as surgical team, wound specialist nurse, dietician, pain team, vascular and diabetes team, podiatrist, physiotherapist, family carers and trained counsellor

## Control or treat underlying causes and barriers to wound healing

Record management plan for: systemic infection, diabetes, nutritional problems, oedema, continence, mobility, vascular issues, pain, stress, anxiety, non-adherence / concordance with offloading and compression, lifestyle choices

## Decide appropriate treatment



\*Use appropriate secondary dressing as per your local protocol; †NPWT: Negative Pressure Wound Therapy; ‡Level of exudate for wounds suitable for NPWT.

## Evaluate and reassess the treatment and wound management outcomes

Evaluate: Record wound progression within given timelines. **Flag** if no change, go back to A, B, C and change treatment where indicated

Developed with the support of Glenn Smith<sup>3</sup> and Moore et al. 2019<sup>4</sup>

§SECURA Range includes SECURA Moisturising Cleanser, SECURA Total Body Foam, SECURA Dimethicone Protectant, SECURA Extra Protective Cream, No Sting Skin Prep; PROSHIELD Range includes PROSHIELD Plus and PROSHIELD Foam and Spray; †ALLEVYN Range includes ALLEVYN LIFE, ALLEVYN GENTLE BORDER and ALLEVYN GENTLE BORDER LITE.

**Reference:** 1. Schultz GS, Sibbald RG, Falanga V, et al. Wound bed preparation: a systematic approach to wound management. *Wound Rep Reg* (2003);11:1-28. 2. Leaper DJ, Schultz G, Carville K, Fletcher J, Swanson T, Drake R. Extending the TIME concept: what have we learned in the past 10 years? *Int Wound J* 2012; 9 (Suppl. 2):1-19. 3. Smith G, Greenwood M, Searle R. Ward nurse's use of wound dressings before and after a bespoke educational programme. *Journal of Wound Care* 2010, vol 19, no.9. 4. Moore Z, Dowsett C, Smith G, et al. TIME CDST: an updated tool to address the current challenges in wound care. *Journal of Wound Care*, vol 28, no 3, March 2019: 154-161.

The products used in the T.I.M.E. clinical decision support tool may vary in different markets. Not all products referred to may be approved for use or available in all markets. Please consult your local Smith+Nephew representative for further details on products available in your market. Intended for healthcare professionals outside of the US only.

Smith+Nephew does not provide medical advice. The information presented is not, and is not intended to serve as, medical advice. For detailed device information, including indications for use, contraindications, precautions and warnings, please consult the product's Instructions for Use (IFU) prior to use. It is the responsibility of healthcare professionals to determine and utilise the appropriate products and techniques according to their own clinical judgment for each of their patients.

Smith+Nephew inc. 2280 Argentia Road, Mississauga, Ont. Canada, L5N-6H8. T 1-800-463-7439 F 1-800-671-9140  
 †Trademark of Smith+Nephew. All Trademarks acknowledged. 13714 | CA53271 04/20