


# Shared Wound Care Mapped



Involve patients with Shared Wound Care

Smith+Nephew



Shaping+  
what's possible  
in wound care

## The path to Shared Wound Care

We know nurses are under more pressure than ever, with chronic wounds outpacing the number of appointments with trained specialist clinicians.<sup>1</sup> Not only that, but patients find dressing change appointments disruptive,<sup>2,3</sup> and up to 30% report depression or anxiety related to their quality of life.<sup>4,5</sup>

By focusing on individual patient needs, we can provide care that works for both patients and practitioners to:

- Develop treatment plans that patients can follow<sup>6-8</sup>
- Improve quality of life and levels of pain, compared to standard care<sup>9,10</sup>
- Increase independence and confidence<sup>8</sup>

With Shared Wound Care we have an opportunity to improve outcomes and empower patients, while also releasing up to 3.5 billion nursing hours globally by 2030.<sup>6-10</sup>

With appropriate clinical tools that help to standardise protocols, reduce variation, and support confident dressing selection, we can shape more empowering wound care experiences for everyone.

Ready to explore your next step?



Indicator at 0%



Indicator 50% Full



Indicator 75% Full

Knowing when to change a dressing can be challenging for both patients and carers. ALLEVYN LIFE<sup>®</sup> Dressing's EXUMASK<sup>®</sup> Change Indicator provides a clear visual guide for timely dressing changes.<sup>15,16</sup>

## I want to find out about the future of Shared Wound Care

Already seeing the benefits of Shared Wound Care and ALLEVYN LIFE Dressings?

Together we can shape what's possible for the future of Shared Wound Care. Sign up for updates.

[Register](#)


## I want a dressing that can increase my confidence in Shared Wound Care

Seeking more confidence in your Shared Wound Care practice?

Discover how ALLEVYN LIFE Dressings deliver trusted performance while saving you valuable time and resources.<sup>11,15,28</sup>

[Download ALLEVYN Dressings brochure](#)

Whether you're a clinician seeking guidance, discussing options with patients, or collaborating with colleagues, we've organised expert-led resources to address your specific Shared Wound Care needs and provide targeted support every step of the way.



## I want to explore Shared Wound Care for my ALLEVYN<sup>®</sup> LIFE Dressing patients

Need guidance on getting started with Shared Wound Care?

The Shared Wound Care Discussion Guide: Made Easy from Wounds International is designed to aid patient engagement and best practice.

[Download Shared Wound Care: Made Easy](#)

## I'm curious about Shared Wound Care

Trying to save time, waste less, and improve lives? Curious about how Shared Wound Care could help your practice?

See how it could empower up to 45% of patients,<sup>7</sup> and save more than 10% of nursing time.<sup>11,13,15,16,27</sup>

[Download Evidence in Focus](#)

Take the next step on your Shared Wound Care path by selecting the stage that best matches your current needs.



## Trusted to work


With unique body-conforming designs, wide borders and flexible layers,<sup>7,11,12</sup> ALLEVYN<sup>®</sup> LIFE Dressings empower patients and help lives to be lived.<sup>13-15</sup>

### EXUMASK<sup>®</sup> change indicator technology

- Minimises the visual impact of absorbed exudate<sup>15,16</sup>
- Works as an indicator as to when to change the dressing which helps minimise clinically unnecessary changes<sup>15,16</sup>

### EXULOCK<sup>®</sup> hyper-absorbent lock-away technology

- Absorbs and locks in exudate to prevent leakage<sup>13,15,17</sup>
- Helps control wound malodour<sup>13-15,17</sup>



Whether it's the extended wear time of dressings,<sup>11,15,18-20</sup> ease of application and removal,<sup>13,21-23</sup> or enabling shared wound care practices,<sup>11,15,24</sup> ALLEVYN products work hard to give you back time.<sup>11,15,21,25,26</sup>

## We know that nurses went into the profession to provide proper care and attention. To make a difference.<sup>29</sup>

Shared Wound Care can make it possible to:



Release more than 10% of nursing hours globally per year<sup>11,13,15,16,27</sup>



Increase collaboration and patient satisfaction, improving nurse-patient relationships<sup>10,30</sup>



Empower patients and enhance independence<sup>6-8</sup>

## Join our community

Register today to stay connected with the Smith+Nephew team and to receive updates about the future of Shared Wound Care

[Register](#)

Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your Smith+Nephew representative or distributor if you have questions about the availability of Smith+Nephew products in your area.

For detailed product information, including indications for use, contraindications, precautions and warnings, please consult the product's applicable Instructions for use (IFU) prior to use.

[www.smith-nephew.com/uk](http://www.smith-nephew.com/uk)

◇Trademark of Smith+Nephew  
All Trademarks acknowledged ©2025 Smith+Nephew 47129-03/25

Smith+Nephew, Croxley Park Building 5  
Hatters Lane, Watford, Hertfordshire,  
WD18 8YE  
**T** +44 (0) 1923 477100  
**F** +44 (0) 1923 477101

### References

1. Moore Z, Probst S. Wounds International. 2023;14(3):28-33. 2. Alam T, Santos V, Woo K. Wounds International. 2018;9(3):6-14. 3. International Consensus. OPTIMISING WELLBEING IN PEOPLE LIVING WITH A WOUND: An Expert Working Group Review; 2012. 4. Renner R, Erfurt-Berge C. Depression and quality of life in patients with chronic wounds: ways to measure their influence and their effect on daily life. Chronic Wound Care Management and Research. 2017;Volume 4:143-151. 5. Olsson M, Järbrink K, Krister, Divakar U, et al. The humanistic and economic burden of chronic wounds: A systematic review. The humanistic and economic burden of chronic wounds. 6. Elsinga B, et al. Wounds International. 2022;London, UK. 7. Moore Z, et al. Wounds International. 2021;12(1):48-53. 8. Loney A, Moore Z. Wounds International. 2023;14(2):1-6. 9. Vahdat S, et al. Patient involvement in health care decision making: A review. Iran Red Crescent Med J. 2014;16(1). 10. Gethin G, et al. J Wound Care. 2020;29(Sup9b):S1-S22. 11. Joy H, Bielby A, Searle R. J Wound Care. 2015;24(7):312, 314-317. 12. Smith+Nephew. Product Performance of Next Generation ALLEVYN Life Internal Report; 2016. 13. Rossington A, et al. Wounds UK. 2013;9(4):91-95. 14. Smith+Nephew. Odour Reducing Properties of ALLEVYN Life. Internal Report; 2012. 15. Tiscar-González V. Adv in Skin Wound Care. 2021;34(1):23-30. 16. Simon D, Bielby A. Wounds UK. 2014;10(3):80-87. 17. Smith+Nephew. Wound Model Testing of New ALLEVYN Life Gen2 Wcl Dressing Using Horse Serum at a Flow Rate Modelling That of a Moderately Exuding Wound; 2016. 18. Smith & Nephew 2017. An Open, Prospective, Randomised, Comparative Volunteer Trial to Compare the Performance of Silicone Adhesive Dressings. Internal Report. GHCA-DOF-10. 19. Smith & Nephew 2017. An Open, Prospective, Randomised, Comparative Study to Compare the Performance of ALLEVYN Gentle Border Multisite with an Alternative Silicone Adhesive Dressing. Internal Report. GHCA-DOF-09. 20. Smith & Nephew 2017. ALLEVYN Gentle Border Wound Model - 10 x 10. Internal Report. DS/17/491/R. 21. Hurd T, et al. Wounds UK. 2009;5(3):32-44. 22. Grothier L, Br J Nurs. 2009;18(11). 23. Callaghan R, et al. Wounds UK. 2012;8(4):110-118. 24. Moore Z, Forni C, McGuinness W. Raising the Standards of care for reducing PU incidence and maintaining skin integrity. In: EPUAP Symposium; 2017. 25. Forni C, Searle R. J Wound Care. 2020;29(2):120-127. 26. Vowden K, Harding K, Moiemien Naiem, Drysdale K, Mistry C. An open, prospective randomised, multi-centre clinical evaluation of a hydrocellular silicone foam dressing\* in the management of exuding chronic and acute wounds. In: Paper Presented at: EWMA; 2011. 27. Moore Z, Loney A, Probst S, et al. Wounds International. 2022;13(2):10-16. 28. Allen D, Saunders C, Tiskowitz K. Reduced weekly dressing changes with a five-layer foam dressing\* compared with other previously used dressings in wounds of mixed aetiology: Results of a systematic literature review and meta-analysis of clinical studies. Paper presented at: SAWC; 2022; Las Vegas. 29. McKenna L. BMC Nurs. 2023;22:152. 30. Loney A, Milne C. Wounds International. 2023;14(2):46-53.