

PICO[◇] 7 and 14 Single Use Negative Pressure Wound Therapy (sNPWT)

Order references

Smith+Nephew

		PICO 7 system		PICO 14 system	Multipack
		1 dressing*	2 dressings†	2 dressings	5 dressings
Dressing	Pad size	S+N code	S+N code	S+N code	S+N code
10cm x 20cm	5.6cm x 15cm	66802012	66802002	66802042	66802022
10cm x 30cm	5.6cm x 25cm	66802013	66802003	66802043	66802023
10cm x 40cm	5.6cm x 35cm	66802014	66802004	66802044	66802024
15cm x 15cm	10cm x 10cm	66802015	66802005	66802045	66802025
15cm x 20cm	10cm x 15cm	66802016	66802006	66802046	66802026
15cm x 30cm	10cm x 25cm	66802017	66802007	66802047	66802027
20cm x 20cm	15cm x 15cm	66802018	66802008	66802048	66802028
25cm x 25cm	20cm x 20cm	66802019	66802009	66802049	66802029
Multisite small 15cm x 20cm	5.5cm x 15cm	66802010	66802000	66802040	66802020
Multisite large 20cm x 25cm	9cm x 20cm	66802011	66802001	66802041	66802021

*1 x dressing kit = 1 dressing + 1 pump †2 x dressing kit = 2 dressings + 1 pump

PICO 7Y system

2 dressings		
Dressing	Pad size	S+N code
Multisite large 20cm x 25cm	9cm x 20cm	66802031

Consumables

Dressing	Description	S+N code
10cm x 12.5cm	Foam wound dressing filler	66801021
11.4cm x 3.7m	5 Antimicrobial Gauze Rolls + 1 SECURA [®] NSBF Wipe	66802127

PICO[®] 7/ 7Y/ 14 sNPWT indications for use:

PICO 7/ 7Y/ 14 sNPWT is indicated for patients who would benefit from a suction device (NPWT) as it may promote wound healing via removal of low to moderate levels of exudate and infectious materials.

Appropriate wound types include: chronic, acute, traumatic, subacute and dehiscent wounds, partial-thickness burns, ulcers (such as diabetic or pressure), flaps and grafts, and closed surgical incisions.

PICO 7/ 7Y/ 14 sNPWT single use negative pressure systems are suitable for use both in a hospital and homecare setting.

PICO 7/ 7Y/ 14 sNPWT is contraindicated for:

- Patients with malignancy in the wound bed or margins of the wound (except in palliative care to enhance quality of life).
- Previously confirmed and untreated osteomyelitis.
- Non-enteric and unexplored fistulas.
- Necrotic tissue with eschar present.
- Exposed arteries, veins, nerves or organs.
- Exposed anastomotic sites

PICO 7/ 7Y/ 14 sNPWT should not be used for the purpose of:

- Emergency airway aspiration.
- Pleural, mediastinal or chest tube drainage.
- Surgical suction.

Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your Smith+Nephew representative or distributor if you have questions about the availability of Smith+Nephew products in your area. For detailed product information, including indications for use, contraindications, precautions and warnings, please consult the product's applicable Instructions for Use (IFU) prior to use.



Smith+Nephew
Croxley Park
Building 5, Lakeside
Hatters Lane, Watford
Hertfordshire WD18 8YE

T: +44 (0) 1923 477100
F: +44 (0) 1923 477101

Smith+Nephew
Kay Fiskers Plads 9
2300 København S
Denmark
T: +45 45 80 61 00

Smith+Nephew
Krokslätts Fabriker 39
431 37 Mölndal
Sweden
T: +46 31 74 65 800

Smith+Nephew
Snarøyveien 36
1364 Fornebu
Norway
T: +47 66 84 20 20

Smith+Nephew
Lentäjantie 1
01530 Vantaa
Finland
T: +358 (0)207 866 33

www.smith-nephew.com

©Trademark of Smith+Nephew
All Trademarks acknowledged
©2024 Smith+Nephew
45067-ukinord 10/24