


PICO[®] sNPWT pathway for non-healing wounds:



For detailed product information, including indications for use, contraindications, precautions and warnings, please consult the product's applicable Instructions for Use (IFU) prior to use

This pathway is adapted from Dowsett, Hampton, et al. 2017,¹ designed to help prevent overuse of the therapy and ensure that patients who may benefit most are able to access the treatment for an optimal period of time.

*Wounds that have healed by <10% but have shown significant improvement in granulation tissue quality/quantity may be considered for further PICO sNPWT treatment based on clinician judgement.

**Wound depth measurements may also be considered as part of the weekly wound assessment. Whether to continue treatment with PICO sNPWT based on changes on wound depth is at the judgment and discretion of the healthcare professional

Smith+Nephew does not provide medical advice. The information presented in this pathway is not, and is not intended to serve as, medical advice. The information presented is for informational purposes only. Only healthcare professionals can advise whether any treatments referred to may be suitable for their patients and specific conditions, and of the associated risks, benefits and possible outcomes. It is the responsibility of healthcare professionals to determine and utilize the appropriate products and techniques according to their own clinical judgment for each of their patients. Individual results will vary.

Reference 1. Dowsett C, Hampton K, Myers D, Styche T. Use of PICO to improve clinical and economic outcomes in hard-to-heal wounds. Wounds International. 2017;8(2):52–58.

Advanced Wound Management. Smith & Nephew, Inc. Fort Worth TX 76109. USA | www.smith-nephew.com | Customer Care Center: 1 800 876-1261; T: 727 392-1261; F: 727 392-6914
 ◊Trademark of Smith+Nephew. All Trademarks acknowledged. ©2025 Smith+Nephew. PCCE38-46520-0725