

Copay Assistance Program:

A few simple facts to get you started

Smith+Nephew

REGRANEX[◇]
(becaplermin) gel 0.01%

Save up to \$200*

Who is eligible to save?*

If you've been prescribed REGRANEX[◇] (becaplermin) gel, 0.01% and pay more than \$25 out of pocket for your prescription, and are commercially insured, you may be eligible to save. **Participants in governmental healthcare programs or where copay assistance is taxed, restricted, or otherwise prohibited by law are not eligible.**

Just how much will I save?

Eligible recipients will pay the first \$25; Smith & Nephew will pay the remaining out-of-pocket cost up to the next \$200.*

How does it work?

The Copay Assistance will be automatically applied at the pharmacy for eligible prescriptions.

-No card necessary.

Please be sure to read the "What is REGRANEX gel?" and "Important Safety Information" sections below, as well as the patient Medication Guide at the end of the REGRANEX gel Prescribing Information.

Copay Assistance Program

Terms and Conditions

Not valid for prescriptions eligible to be reimbursed under Medicare (including Medicare part D and Medicare Advantage), Medicaid, TRICARE, CHAMPUS, the Puerto Rico Governmental Health Insurance Plan, or other federal, state, or governmental healthcare programs, or uninsured patients. Valid in the U.S. only, void where taxed, restricted, or prohibited by law. Accepted by participating pharmacies only. This Copay Assistance Program is not insurance and may be changed or discontinued at any time without notice.





What is REGRANEX Gel?

REGRANEX gel is a protein medicine made in yeast that is used with other ulcer care practices (such as good wound care) to treat diabetic sores (ulcers) of your legs or feet that are deeper than just your skin, in people who have good blood supply to the legs.

It is not known if REGRANEX gel is effective for the treatment of pressure ulcers or ulcers that are due to poor blood flow (circulation). It is not known if REGRANEX gel is safe and effective in children under 16 years of age.

Important Safety Information

REGRANEX (becaplermin) gel 0.01% is indicated for treating lower extremity diabetic ulcers that extend under the skin and have adequate blood supply. The effectiveness of REGRANEX gel has not been established in pressure ulcers, or venous ulcers. REGRANEX gel is not substitute for good ulcer care practices. REGRANEX gel is contraindicated for patients with a tumor at the site(s) of application. REGRANEX gel is non-sterile so it should not be used in wounds that heal without tissue loss (primary intention). Side effects include rashes, burning sensation, and redness at the site of application. Tumors distant from the site of application have been reported in both a clinical study and in real world use. The risk information provided herein is not comprehensive. To see the complete prescribing information, please see the FDA-approved product labeling, here: https://regranex.com/pdf/PI_Full_Version.pdf. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch or call 1-800-FDA-1088.

www.smith-nephew.com

This website is intended for individuals age 18 and over in the United States only. The information provided is for information purposes only and is not intended to be medical advice. It does not replace the professional training, experience, and knowledge of the healthcare provider responsible for patient care, who must base treatment upon the unique characteristics of each patient. If you have questions about REGRANEX (becaplermin) gel, 0.01% that are not answered by this website; contact your healthcare provider.

Advanced Wound Management
Smith & Nephew, Inc.
Fort Worth, TX 76109 USA

Customer Care Center
1-800-876-1261
T 727-392-1261
F 727-392-6914

◇Trademark of Smith+Nephew
©2023 Smith+Nephew, Inc.
RGWE15-38461-0124