Breaking down barriers

Progressing wounds

SmithNephew

IODOSORB° gel and IODOFLEX° dressing is indicated for use in chronic and infected wounds where bioburden is a barrier to healing.

Prepare the wound to progress to closure.





 IODOSORB gel has been shown to physically disrupt mature biofilm by absorbing it into the dressing and reducing the bacterial load within the biofilm.^{11-17*}

*Across (in-vitro)and animal models



- Provides broad-spectrum antimicrobial activity (in vitro) for up to 72 hours even when re-challenged daily^{18,19}
- More effective in reducing bioburden and infection compared to standard care²⁰




DEBRIS AND BACTERIA

2-720 22-24

- Provides rapid and effective desloughing^{2-7,20,23-24}
- Debrides wound bed via absorption of excess slough and debris to prepare for effective healing^{11,13,20,23}

IODOSORB° gel and IODOFLEX° dressing unique antimicrobial dressing made of cadexomer micro-beads: spherical starch structures loaded with 0.9% elemental iodine.



Ordering codes for IODOSORB gel and IODOFLEX dressing				
IODOSORB Gel 0.9% Cadexomer Iodine	Size	Product #	Pcs/Pkg	Pkgs/Unit
	40g	6602125040	1	12
	10g	6602124014	4	12
IODOFLEX Pads 0.9% Cadexomer Iodine	5g (1½" x 2¾")	6602133005	5	12
	10g (2³¼" x 3")	6602134010	3	12

For detailed product information on indication, contraindication and important safety information, please see the Instructions for Use (IFU).

Advanced Wound Management www.smith-nephew.com Smith & Nephew

Fort Worth, TX 76109 USA

1800876-1261 T 727 392-1261 F727392-6914 *Trademark of Smith & Nephew ©2022 Smith & Nephew IOPE17-32565-1022

References: 1. Skog E, et al. A randomized trial comparing cadexomer iodine and standard treatment in the out-patient management of chronic venous ulcers. Br. J. Dermatol. 109, 77–83 (1983), 2. Troëng T, Skog E, Arnesjö B, Gjöres JE, Bergljung L, Gundersen JEA. A randomised multicentre trial to compare the efficacy of cadexomer iodine and standard treatment in the management of chronic venous ulcers in out patients. in Cadexomer Iodine, (eds. Fox, J. & Fisher, H.) 43-50 (Schattauer Verlag, 1983). 3. Ormiston MC, Fox JA. Controlled trial of Iodósorb. Br. Med. J. (Clin. Res. Ed). 291, 308-310 (1985) 4. Hansson C, et al. The effects of cadexomer iodine paste in the treatment of venous leg ulcers compared with hydrocolloid dressing and paraffin gauze dressing. Int. J. Dermatol. 37, 390–396 (1998). 5. Lindsay G, Latta D, Lyons KGB. A study in general practice of the efficacy of cadexomer iodine in venous leg ulcers treated on alternate days. Acta Ther. 12, 141-148 (1986). 6. Johnson A. A combative healer with no ill-effect. Iodosorb in the treatment of infected wounds. Prof. Nurse 7, 60, 62, 64 (1991). 7. Harcup JW, Saul PA. A study of the effect of cadexomer iodine in the treatment of venous leg ulcers. Br. J. Clin. Pract. 40, 360-4 (1986). 8. Hesler B. IODOSORB[™] Powder wound model testing. 3 day cavity testing with ALLEVYN[™] GENTLE Border secondary dressing using horse serum at a flow rate modelling that of a moderately exuding wound. Smith&Nephew Data on File # DS.17.249.R. (2017). 9. Hesler B. IODOSORB Ointment/Gel wound model testing. 3 day cavity testing with ALLEVYN gentle border secondary dressing using horse serum at a flow rate modelling that of a moderately exuding wound. Smith&Nephew Data on File DS.17.572.R. (2017). 10. Hesler B. IODOSORB IODOFLEX™ dressing wound model testing. 3 day cavity testing with ALLEVYN gentle border secondary dressing using horse serum at a flow rate modelling that of a moderately exuding wound. Smith&Nephew Data on File # DS.17.247.R. (2017). 11. Fitzgerald, D. J. et al.. Wound Repair Regen. 25, 13–24 (2017), 12. Akiyama, H., Oono, T., Saito, M. & Iwatsuki, K. J. Dermatol. 31, 529-34 (2004), 13. Roche, E. et al. Int. Wound J. In press, 1-10 (2019), 14. Oates, J. L., Phillips, C. D., Wolcott, R. & Woodmansey, E. in SAWC Atlanta 1 (2016), 15. Phillips, P. L. et al.. Int. Wound J. 469-483 (2013), 16. Hill, K. E. et al. J. Antimicrob. Chemother. 65, 1195-1206 (2010), 17. Kiamco, M. M., Atci, E., Mohamed, A., Call, D. R. & Beyenal, H. Appl. Environ. Microbiol. 1-33 (2017) 18. Forest E. Antimicrobial activity of IODOSORB range against a broad spectrum of wound pathogens. Smith&Nephew Data on File #1801001. (2018). 19. Harrow J. A comparison of the antimicrobial activity of a cadexomer iodine dressing and a povidone iodine dressing. 6763/IODOSORB/TECHMON/GLOBAL/0404. (2009). 20. Sundberg J, Pharm MS, Meller R, Se B, Pharm MS. A retrospective Review of the Use of Cadexomer Iodine in the Treatment of Chronic Wounds. Wounds A Compend. Clin. Res. Pract. 9, 68–86 (1997). 21. Akiyama H, Oono T, Saito M, Iwatsuki K. Assessment of cadexomer iodine against Staphylococcus aureus biofilm in vivo and in vitro using confocal laser scanning microscopy. J. Dermatol. 31, 529–34 (2004). 22. Zhou LH, Nahm WK, Badiavas E, Yufit T, Falanga V. Slow release iodine preparation and wound healing: in vitro effects consistent with lack of in vivo toxicity in human chronic wounds. Br. J. Dermatol. 146, 365-74 (2002). 23. Holloway GÁ, Johansen KH, Barnes RW, Pierce GE. Multicenter trial of cadexomer iodine to treat venous stasis ulcer. West. J. Med. 151, 35–38 (1989). 24. Moberg S, Hoffman MOL, Holst A. A Randomized Trial of Cadexomer in Decubitus Ulcers Iodine. 25. Laudanska H, Gustavson B. In-patient treatment of chronic varicose venous ulcers. A randomized trial of cadexomer iodine versus standard dressings. J. Int. Med. Res. 16, 428–35 (1988) 26. Hesler B. Release of iodine from IODOSORB/IODOFLEX dressing into 0.9% saline solution over a three day period. Smith&Nephew Data on File # DS.18.026.R. (2018). 27. Hesler B. Release of iodine from IODOSORB powder into 0.9% saline solution over a three day period. Smith&Nephew Data on File # DS.18.024.R. (2018). 28. Hesler B. Release of iodine from IODOSORB ointment into 0.9% saline solution over a three day period. Smith&Nephew Data on File # DS.18.025.R. (2018). 29. Moberg S, Hoffman L, Grennert ML, Holst A. A randomized trial of cadexomer iodine in decubitus ulcers. J. Am. Geriatr. Soc. 31, 462-465 (1983). 30. Hesler B. IODOSORB Powder wound model testing to investigate colour loss of the product over the course of a 3 day period. Data on file # DS.17.594.R. (2017). 32. Hesler B. (2017) Method transfer report Smith+Nephew Data on File # DS.17.210.R.,