Test Results of the Effects of Silver Antimicrobial Dressings on Drug Resistant Bacteria *in vitro*

SmithNephew

ACTICOAT

Antimicrobial Barrier Dressings

Brand name	Product types available	Log reduction in 30 minutes for MRSA	Log reduction in 30 minutes for VRE	Log reduction in 30 minutes for 5 NDM-1 producing bacterial strains	Log reduction in 30 minutes for drug resistant P. aeruginosa 137366	Method of silver delivery		What does that mean?
						Initial	Solution (concentration)	What does that mean?
ACTICOAT° Antimicrobial Barrier Dressing (Smith+Nephew)	Dressing (3 and 7 day)	>6.01	>6.01	>4.07	>5.08	Silver Ag ^o Nanocrystals ¹⁰	Ag+, Ag ⁰ / Ag+ clusters (70-100 ppm) ¹⁰	ACTICOAT dressng showed marked bactericidal activity against MRSA, VRE, NDM-1 producing bacterial strains and drug-resistant P. aeruginosa at 30 minutes (<i>in vitro</i>)
Silverlon™ (Argentum LLC)	Dressing, island dressing,foam strips (7 day)	0.01	1.541	Not reported	<18	Silver oxide ¹¹	Ag+ (<1 ppm)11	Silverlon showed no activity against MRSA and bacteriostatic activity against VRE within 30 minutes. No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drugresistant P. aeruginosa
AQUACEL™Ag (ConvaTec)	Dressing, rope (7 Day)	0.01	0.01	Not reported	<18	AgCl ¹¹	Ag+ (<1-3 ppm) ¹ _a	Aquacel Ag showed no activity against either MRSA or VRE within 30 minutes. No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drug-resistant P. aeruginosa
Aquacel™ Ag ExtraTM (ConvaTec)	Dressing (7 Days)	<12	<12	Not reported	0.02	AgCI ¹¹	Ag+ (<1-3 ppm) ¹	Aquacel Ag Extra showed no activity against either MRSA or VRE within 30 minutes. No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drug-resistant P. aeruginosa
TheraBond 3D™	Wound layer, island, wrap dressing(7 day)	0.03	No Speed of kill data	Not reported	<13	Silver plated fabric	Ag+ Not reported	Therabond 3D demonstrated bactericidal activity against MRSA, but no rapid activity against VRE was reported. No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drugresistant P. aeruginosa
KerraContact™ (Crawford Healthcare)	Barrier dressing (7 day)	0.04	0.04	Not reported	Not reported	Silveroxysalts ¹²	Ag+, Ag+2, Ag3+ (6 ppm) ^{12,13}	KerraContact is bactericidal against MRSA or VRE from 4 hours with no effect at 30 minutes reported. No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drug-resistant P. aeruginosa
Mepilex [™] Ag (Mölnlycke)	Foam, post-op border, transfer (7 day)	<1 ⁵	<16	Not reported	<19	Silver sulphate ¹⁴	Ag+ (6 ppm) ^{15,16}	Mepilex Ag showed no activity against MRSA or VRE in 30 minutes. No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drug-resistant P. aeruginosa
Silvercel™ (Systagenix)	Dressing, rope (7 day)	<16	0.06	Not reported	Not reported	Ag alginate ¹⁷	Ag+ (~1 ppm) ¹¹	Silvercel showed no activity against MRSA or VRE in 30 minutes. No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drug-resistant P. aeruginosa
SilvaSorb™ (Medline Ind.)	Sheet, perforated sheet, strands, (7 day), gel(3 day)	0.01	0.01	Not reported	<18	AgCl11	Ag+ (~1 ppm) ¹¹	SilvaSorb showed no activity against MRSA or VRE in 30 minutes No published data to support rapid antimicrobial activity against NDM-1 producing bacterial strains or drug-resistant P. aeruginosa

The Minimum Bactericidal Concentration (MBC) for clinically relevant bacteria is between 5 and >50ppm^{18,19} and up to 60.5ppm for MRSA20. This means in order to kill MRSA the dressing must kill bacteria with a MBC of at least 60.5 PPM (200ppm x 0.0325=60.5ppm). Notes: In vitro results do not necessarily translate into clinically relevant results. Antimicrobial agents are considered bactericidal if they induce a 3 or greater log reduction in a clinically relevant timeframe¹⁹ These results do not represent a head-to-head safety/efficacy clinical comparison.

Test Results of the Effects of Silver Antimicrobial Dressings on Drug Resistant Bacteria *in vitro*



ACTICOAT[♦]
Antimicrobial Barrier Dressings

For detailed product information, including indications for use, contraindications, effects, warnings and precautions, please consult the product's Instructions for Use (IFU) prior to use.

References

- 1. Data on file: Testing performed by Hoosier Microbiological Laboratory, Muncie, IN (independent lab). January 3, 2005.
- 2. Data on file: 1405007. Antimicrobial activity of DURAFIBER™, DURAFIBER Ag, Aquacel™ Extra, Aquacel Ag Extra and Aquacel Ag+ Extra against bacteria, yeast and fungi commonly found in wounds over a 7 day period.
- 3. Data on file: 0911020. Antimicrobial assessment of ACTICOAT™ Flex 7 in comparison to Therabond™ against MRSA and Pseudomonas aeruginosa, 2009.
- Exsalt[™] SD7-4 Brochure. PR-0045.0.5.Data on file: 070805
- 5. Antimicrobial activity of ALLEVYN™ Ag Dressings in comparison to other silver-based antimicrobial dressings against MRSA using a dynamicshake flask method, 2007.
- 6. Data on file: Testing performed by Hoosier Microbiological Laboratory, Muncie, IN (independent lab). Raw data/XL file.
- 7. Hope R, et al. The in vitro antibacterial activity of Nanocrystalline Silver dressings against bacteria with NDM-1 carbapenemase [poster]. EWMA; 2012.
- 8. Data on file: Testing performed by Hoosier Microbiological Laboratory, Muncie, IN (independent lab). January 4, 2005.
- Data on file: 0707051. Antimicrobial activity of ALLEVYN Ag Dressings in comparison to other silver-based antimicrobial dressings against Pseudomonas aeruginosausing a
 dynamic shake flask method, 2007.
- 10. Dunn K, et al. The role of ACTICOAT with Nanocrystalline Silver in the management of burns. Burns. 2004; 30 Suppl 1:S1-9.
- 11. Ovington LG. Why is Nanocrystalline Silver Superior? WOUNDS. 2004; 13(2):1-9.
- 12. Exsalt SD7-4 Brochure. PR-0046.0.
- 13. Djokić N, et al. Coating of High Density Polyethylene (HDPE) with Silver Compounds and Their Antimicrobial Activity in vitro. ECS Transactions. 2008; 11(21):59-68.
- 14. Mepilex[™] Ag 510k (K071354).
- 15. Schwarzkopf A, et al. "The Release of Silver Ions from Foam Dressings". Wound Management. 2010; 4(1): 1-4.
- 16. Mepilex Ag European Patent Application: 07004275.9.
- 17. Silvercel™ Brochure, AWC-SC001.
- 18. Yin, H.Q. et al. Comparative Evaluation of the Antimicrobial Activity of ACTICOAT Antimicrobial Barrier Dressing. J Burn Care Rehabil 1999;20:195-200.
- 19. Hall, R.E. et al. Inhibitory and Cidal Antimicrobial Actions of Electrically Generated Silver Ions. J Oral Maxillofac Surg 1987;45:779-784.
- 20. Maple, P.A. et al. Comparison of the in vitro activities of the topical antimicrobials azelaic acid, nitrofurazone, silver sulphadiazine and mupirocin against methicillin-resistant Staphylococcus aureus. Journal of Antimicrobial Chemotherapy. 1992;29:661-668.

Advanced Wound Management Smith & Nephew Fort Worth, TX 76109 USA T +1 (800) 876 1261

F+1 (727) 392 6914

◇Trademark of Smith+Nephew ©2022 Smith+Nephew ACPE8-32565-1022

www.smith-nephew.com