
+ Case study

SmithNephew

Use of ACTICOAT FLEX 3 Antimicrobial Barrier Dressing to help prevent infection in bilateral lower limb superficial burns

Presentation

At the clinic, a 50-year-old female, with no known comorbidities and whose activities of daily living include full-time care of dependents, presented with one-week old superficial burns on both lower limbs (total area: $75 \times 20 \times 0.25$ cm) caused by hot water. The burns were restricting her mobility (bending of the knees) and negatively impacting her self-esteem.



Treatment

When the burn first occurred, the patient was treated with a hydrogel dressing at the emergency room. Mobility was encouraged at all visits.

Visit 1: Start of treatment







The patient was seen by a nurse specialist at the wound clinic one week later who was concerned about the potential for infection and immobility related to scar tissue. ACTICOAT FLEX 3 Dressing was applied with an absorbent dressing held in place using crepe bandaging.

Visit 2: Day 3: Follow-up treatment visit







The wounds had reduced in size (total area 60 x 16 x 0cm) and ACTICOAT FLEX 3 Dressings were reapplied.

+ Case study

Smith-Nephew

Visit 3: Day 6: Completion of treatment



The wounds had epithelialised and the patient was advised to apply emollient moisturiser twice daily to the area.



ACTICOAT FLEX 3 Antimicrobial Barrier Dressing

Outcomes





The patient commented on **relief** from pain and inflammation regained full mobility allowing her to resume daily activities within two weeks of occurrence



Inflammation and oedema were notably reduced after application nurse to result in **faster healing**



easy to use and conformable

Summary

- Following treatment, including ACTICOAT FLEX 3 Dressing:
 - The patient was fully mobile and healed after 2 weeks, without developing signs or symptoms of infection
 - Returned to her usual routine within 2 weeks, preventing disruption of the family unit
- · ACTICOAT FLEX 3 Dressing was reported by the specialist nurse to help reduce inflammation and oedema, thereby promoting healing, while being easy to use and conformable

Acknowledgements: With kind permission of M Second, Specialist Nurse, Waterfall City Hospital, Midrand, South Africa.